

Elimination of quantifiers in the theory of projectable real closed rings with the first convexity property.

Jorge I. Guier – Escuela de Matemática y Centro de Investigaciones en Matemática Pura y Aplicada (CIMPA) – Universidad de Costa Rica

August 8, Banyuls-sur-Mer, DDG40, 2025.

First notions

An f -ring is a subdirect products of totally ordered rings (a lattice ordered ring). This is expressible by an universal first order sentence in the language of lattice-ordered rings $\mathcal{L}_{\text{lor}} = \{0, 1, +, -, \cdot, \wedge\}$. Precisely:

$$\forall a \forall b \forall x (a \wedge 0 = b \wedge 0 = x \wedge 0 = 0 \wedge a \wedge b = 0 \rightarrow a \wedge bx = a \wedge xb = 0).$$

First notions

An f -ring is a subdirect products of totally ordered rings (a lattice ordered ring). This is expressible by an universal first order sentence in the language of lattice-ordered rings $\mathcal{L}_{\text{lor}} = \{0, 1, +, -, \cdot, \wedge\}$. Precisely:

$$\forall a \forall b \forall x (a \wedge 0 = b \wedge 0 = x \wedge 0 = 0 \wedge a \wedge b = 0 \rightarrow a \wedge bx = a \wedge xb = 0).$$

The language of ordered rings will be denoted by $\mathcal{L}_{\text{or}} = \{0, 1, +, -, \cdot, <\}$.

First notions

An f -ring is a subdirect products of totally ordered rings (a lattice ordered ring). This is expressible by an universal first order sentence in the language of lattice-ordered rings $\mathcal{L}_{\text{lor}} = \{0, 1, +, -, \cdot, \wedge\}$. Precisely:

$$\forall a \forall b \forall x (a \wedge 0 = b \wedge 0 = x \wedge 0 = 0 \wedge a \wedge b = 0 \rightarrow a \wedge bx = a \wedge xb = 0).$$

The language of ordered rings will be denoted by $\mathcal{L}_{\text{or}} = \{0, 1, +, -, \cdot, <\}$.

An f -ring A is **projectable** if $A = a^\perp + a^{\perp\perp}$, for all $a \in A$; where the **polar** of $a \in A$ is $a^\perp = \{b \in A : a \perp b\}$ and the **bipolar** of a is $a^{\perp\perp} = \{b \in A : b \perp c \text{ for all } c \in a^\perp\}$. This notion is expressed by a first order formula in $\mathcal{L}_{\text{lor}}$.

First notions

An f -ring is a subdirect products of totally ordered rings (a lattice ordered ring). This is expressible by an universal first order sentence in the language of lattice-ordered rings $\mathcal{L}_{\text{lor}} = \{0, 1, +, -, \cdot, \wedge\}$. Precisely:

$$\forall a \forall b \forall x (a \wedge 0 = b \wedge 0 = x \wedge 0 = 0 \wedge a \wedge b = 0 \rightarrow a \wedge bx = a \wedge xb = 0).$$

The language of ordered rings will be denoted by $\mathcal{L}_{\text{or}} = \{0, 1, +, -, \cdot, <\}$.

An f -ring A is **projectable** if $A = a^\perp + a^{\perp\perp}$, for all $a \in A$; where the **polar** of $a \in A$ is $a^\perp = \{b \in A : a \perp b\}$ and the **bipolar** of a is $a^{\perp\perp} = \{b \in A : b \perp c \text{ for all } c \in a^\perp\}$. This notion is expressed by a first order formula in $\mathcal{L}_{\text{lor}}$.

A ring is **reduced** if it doesn't have nilpotent elements other than zero.

First notions

An f -ring is a subdirect products of totally ordered rings (a lattice ordered ring). This is expressible by an universal first order sentence in the language of lattice-ordered rings $\mathcal{L}_{\text{lor}} = \{0, 1, +, -, \cdot, \wedge\}$. Precisely:

$$\forall a \forall b \forall x (a \wedge 0 = b \wedge 0 = x \wedge 0 = 0 \wedge a \wedge b = 0 \rightarrow a \wedge bx = a \wedge xb = 0).$$

The language of ordered rings will be denoted by $\mathcal{L}_{\text{or}} = \{0, 1, +, -, \cdot, <\}$.

An f -ring A is **projectable** if $A = a^\perp + a^{\perp\perp}$, for all $a \in A$; where the **polar** of $a \in A$ is $a^\perp = \{b \in A : a \perp b\}$ and the **bipolar** of a is $a^{\perp\perp} = \{b \in A : b \perp c \text{ for all } c \in a^\perp\}$. This notion is expressed by a first order formula in $\mathcal{L}_{\text{lor}}$.

A ring is **reduced** if it doesn't have nilpotent elements other than zero.

An ordered ring A satisfies the **first convexity property** if

$$\forall a \forall b (0 < a < b \rightarrow b \mid a),$$

is true in A .

First notions

A lattice ordered ring is called **divisible-projectable** if it satisfies the following formula:

$$\forall x \forall y \left(y \neq 0 \rightarrow \exists z \exists w (x = z + w \wedge z \perp w \wedge y \mid z \wedge$$

$$\forall w' (w' \neq 0 \wedge w' \perp (w - w') \rightarrow y \nmid w')) \right).$$

First notions

A lattice ordered ring is called **divisible-projectable** if it satisfies the following formula:

$$\forall x \forall y \left(y \neq 0 \rightarrow \exists z \exists w (x = z + w \wedge z \perp w \wedge y \mid z \wedge$$

$$\forall w' (w' \neq 0 \wedge w' \perp (w - w') \rightarrow y \nmid w')) \right).$$

A ring A is called **sc-regular** if there exists an element $u \in A$ such that $\text{Ann}(u) = \{0\}$ (or $1 \preceq u$) and $u \nmid e$ for every non-zero idempotent $e \in A$.

First notions

A lattice ordered ring is called **divisible-projectable** if it satisfies the following formula:

$$\forall x \forall y \left(y \neq 0 \rightarrow \exists z \exists w (x = z + w \wedge z \perp w \wedge y \mid z \wedge$$

$$\forall w' (w' \neq 0 \wedge w' \perp (w - w') \rightarrow y \nmid w')) \right).$$

A ring A is called **sc-regular** if there exists an element $u \in A$ such that $\text{Ann}(u) = \{0\}$ (or $1 \preceq u$) and $u \nmid e$ for every non-zero idempotent $e \in A$.

Real closed rings have been introduced by Niels Schwartz in the late 80's. There are many different ways to see those rings (isomorphic to its real closure, an inductive definition using first the notion of a real closed domain, there is an axiomatization, etc).

First notions

A lattice ordered ring is called **divisible-projectable** if it satisfies the following formula:

$$\forall x \forall y \left(y \neq 0 \rightarrow \exists z \exists w (x = z + w \wedge z \perp w \wedge y \mid z \wedge$$

$$\forall w' (w' \neq 0 \wedge w' \perp (w - w') \rightarrow y \nmid w')) \right).$$

A ring A is called **sc-regular** if there exists an element $u \in A$ such that $\text{Ann}(u) = \{0\}$ (or $1 \preceq u$) and $u \nmid e$ for every non-zero idempotent $e \in A$.

Real closed rings have been introduced by Niels Schwartz in the late 80's. There are many different ways to see those rings (isomorphic to its real closure, an inductive definition using first the notion of a real closed domain, there is an axiomatization, etc).

In the context of this talk, you may think real closedness equivalent to the satisfaction of the intermediate value property for polynomials.

Cherlin-Dickmann theorem

Cherlin and Dickmann proved (1983) that the theory of **real closed valuation rings** admits elimination of quantifiers in the language $\mathcal{L}_{\text{or}} \cup \{|\}$, where $|$ is the divisibility relation.

Boolean Products

Let L be a first-order language, $\mathfrak{A}$ a L -structure and $\{\mathfrak{A}_x : x \in X\}$ a family of L -structures.

Boolean Products

Let L be a first-order language, $\mathfrak{A}$ a L -structure and $\{\mathfrak{A}_x : x \in X\}$ a family of L -structures.

We say that $\mathfrak{A}$ is a **Boolean product** of $\{\mathfrak{A}_x : x \in X\}$ in L , denoted by $\mathfrak{A} \in \Gamma_L^{\mathfrak{A}}(X, (\mathfrak{A}_x)_{x \in X})$, if the following conditions holds:

Boolean Products

Let L be a first-order language, $\mathfrak{A}$ a L -structure and $\{\mathfrak{A}_x : x \in X\}$ a family of L -structures.

We say that $\mathfrak{A}$ is a **Boolean product** of $\{\mathfrak{A}_x : x \in X\}$ in L , denoted by $\mathfrak{A} \in \Gamma_L^{\mathfrak{A}}(X, (\mathfrak{A}_x)_{x \in X})$, if the following conditions holds:

(i) X is a Boolean space.

Boolean Products

Let L be a first-order language, $\mathfrak{A}$ a L -structure and $\{\mathfrak{A}_x : x \in X\}$ a family of L -structures.

We say that $\mathfrak{A}$ is a **Boolean product** of $\{\mathfrak{A}_x : x \in X\}$ in L , denoted by $\mathfrak{A} \in \Gamma_L^{\mathfrak{A}}(X, (\mathfrak{A}_x)_{x \in X})$, if the following conditions holds:

- (i) X is a Boolean space.
- (ii) $\mathfrak{A}$ is a subdirect product of $\{\mathfrak{A}_x : x \in X\}$.

Boolean Products

Let L be a first-order language, $\mathfrak{A}$ a L -structure and $\{\mathfrak{A}_x : x \in X\}$ a family of L -structures.

We say that $\mathfrak{A}$ is a **Boolean product** of $\{\mathfrak{A}_x : x \in X\}$ in L , denoted by $\mathfrak{A} \in \Gamma_L^{\mathfrak{A}}(X, (\mathfrak{A}_x)_{x \in X})$, if the following conditions holds:

- (i) X is a Boolean space.
- (ii) $\mathfrak{A}$ is a subdirect product of $\{\mathfrak{A}_x : x \in X\}$.
- (iii) For every atomic L -formula $\Phi(v_1, \dots, v_n)$ and every $a_1, \dots, a_n \in |\mathfrak{A}|$,

$$\llbracket \Phi(a_1, \dots, a_n) \rrbracket =_{\text{def}} \{x \in X : \mathfrak{A}_x \models \Phi(a_1(x), \dots, a_n(x))\}$$

is a clopen subset of X .

Boolean Products

Let L be a first-order language, $\mathfrak{A}$ a L -structure and $\{\mathfrak{A}_x : x \in X\}$ a family of L -structures.

We say that $\mathfrak{A}$ is a **Boolean product** of $\{\mathfrak{A}_x : x \in X\}$ in L , denoted by $\mathfrak{A} \in \Gamma_L^{\mathfrak{A}}(X, (\mathfrak{A}_x)_{x \in X})$, if the following conditions holds:

- (i) X is a Boolean space.
- (ii) $\mathfrak{A}$ is a subdirect product of $\{\mathfrak{A}_x : x \in X\}$.
- (iii) For every atomic L -formula $\Phi(v_1, \dots, v_n)$ and every $a_1, \dots, a_n \in |\mathfrak{A}|$,

$$\llbracket \Phi(a_1, \dots, a_n) \rrbracket =_{\text{def}} \{x \in X : \mathfrak{A}_x \models \Phi(a_1(x), \dots, a_n(x))\}$$

is a clopen subset of X .

- (iv) Patchwork property: For every $a, b \in \mathfrak{A}$ and any clopen set N of X , the element $c = a \upharpoonright_N \cup b \upharpoonright_{X \setminus N}$ defined by

$$c(y) = \begin{cases} a(y) & \text{if } y \in N \\ b(y) & \text{if } y \in X \setminus N, \end{cases}$$

belongs to $|\mathfrak{A}|$.

Boolean Products

We say that $\mathfrak{A}$ is an **elementary Boolean product** of $\{\mathfrak{A}_x : x \in X\}$ in L , denoted by $\mathfrak{A} \in \Gamma_L^e(X, (\mathfrak{A}_x)_{x \in X})$, if $\mathfrak{A}$ is a Boolean product of $\{\mathfrak{A}_x : x \in X\}$ in L and condition (iii) is verified for **all** L -formulas $\Phi(v_1, \dots, v_n)$.

Boolean Products

We say that $\mathfrak{A}$ is an **elementary Boolean product** of $\{\mathfrak{A}_x : x \in X\}$ in L , denoted by $\mathfrak{A} \in \Gamma_L^e(X, (\mathfrak{A}_x)_{x \in X})$, if $\mathfrak{A}$ is a Boolean product of $\{\mathfrak{A}_x : x \in X\}$ in L and condition (iii) is verified for **all** L -formulas $\Phi(v_1, \dots, v_n)$.

Let A be a reduced f -ring. Then A is a projectable, divisible-projectable, sc-regular real closed ring with the first convexity property if and only if A is the ring of continuous sections of a Hausdorff sheaf of Real Closed Valuation Rings (in the language of ordered rings adjoining the divisibility relation), that is:

$$A \in \Gamma_{\mathcal{L}_{\text{or}} \cup \{|\cdot|\}}^e(\pi A, (A/p)_{p \in \pi A}),$$

where πA is the space of minimal prime ideals of A and A/p is a real closed valuation ring, for every $p \in \pi A$.

Radical relations were defined by Prestel-Schwartz in the following way:

- (1) $a \preceq a$, for all $a \in A$;
- (2) if $a \preceq b$ and $b \preceq c$ then $a \preceq c$, for all $a, b, c \in A$;
- (3) if $a \preceq c$ and $b \preceq c$ then $a + b \preceq c$, for all $a, b, c \in A$;
- (4) if $a \preceq b$ then $ac \preceq bc$, for all $a, b, c \in A$;
- (5) $a \preceq 1$, for all $a \in A$ and $1 \not\preceq 0$;
- (6) $b \preceq b^2$, for all $b \in A$.

Radical relations and von Neumann regular rings

Prestel-Schmid (1990) showed that for any radical relation $\preceq$ on a ring A , there exists a subset $X \subseteq \operatorname{Spec}(A)$ such that:

$$a \preceq b \iff \forall p \in X (a \notin p \Rightarrow b \notin p).$$

Radical relations and von Neumann regular rings

Prestel-Schmid (1990) showed that for any radical relation $\preceq$ on a ring A , there exists a subset $X \subseteq \text{Spec}(A)$ such that:

$$a \preceq b \iff \forall p \in X (a \notin p \Rightarrow b \notin p).$$

Prestel-Schwartz (2002) showed:

- if $X = \pi A$ is the space of minimal prime ideals of a ring A , then the radical relation given by X is

$$a \preceq_{\pi A} b \iff \forall x (bx = 0 \rightarrow ax = 0) \iff \text{Ann}(b) \subseteq \text{Ann}(a).$$

Radical relations and von Neumann regular rings

Prestel-Schmid (1990) showed that for any radical relation $\preceq$ on a ring A , there exists a subset $X \subseteq \text{Spec}(A)$ such that:

$$a \preceq b \iff \forall p \in X (a \notin p \Rightarrow b \notin p).$$

Prestel-Schwartz (2002) showed:

- if $X = \pi A$ is the space of minimal prime ideals of a ring A , then the radical relation given by X is

$$a \preceq_{\pi A} b \iff \forall x (bx = 0 \rightarrow ax = 0) \iff \text{Ann}(b) \subseteq \text{Ann}(a).$$

- the theory of von Neumann regular real closed rings without non-zero minimal idempotents admits elimination of quantifiers in the language $\mathcal{L}_{\text{lor}} \cup \{\preceq\}$.

A theory of real closed rings and Local Divisibility

Let T^* be the theory of projectable, sc-regular, divisible-projectable, real closed rings without minimal (non-zero) idempotents.

A theory of real closed rings and Local Divisibility

Let T^* be the theory of projectable, sc-regular, divisible-projectable, real closed rings without minimal (non-zero) idempotents.

The local divisibility relation is defined by:

$$y \mid_{\text{loc}} w \longleftrightarrow \exists w' (w' \neq 0 \wedge w'(w - w') = 0 \wedge y \mid w') \vee w = 0,$$

or equivalently:

$$y \mid_{\text{loc}} w \longleftrightarrow \exists e [e^2 = e \wedge we \neq 0 \wedge e \preceq w \wedge y \mid we] \vee w = 0.$$

A theory of real closed rings and Local Divisibility

Let T^* be the theory of projectable, sc-regular, divisible-projectable, real closed rings without minimal (non-zero) idempotents.

The local divisibility relation is defined by:

$$y \mid_{\text{loc}} w \longleftrightarrow \exists w' (w' \neq 0 \wedge w'(w - w') = 0 \wedge y \mid w') \vee w = 0,$$

or equivalently:

$$y \mid_{\text{loc}} w \longleftrightarrow \exists e [e^2 = e \wedge we \neq 0 \wedge e \preceq w \wedge y \mid we] \vee w = 0.$$

Theorem (G., 2025): The theory T^* admits elimination of quantifiers in the language $\mathcal{L}_{\text{lor}} \cup \{|\, , \preceq, \mid_{\text{loc}}\}$.

Idea of the proof

The model completeness of the theory T^* is established in the language of $\mathcal{L}_{\text{lor}} \cup \{\preceq, |_{\text{loc}}\}$ using Comer's result (a Feferman-Vaught result for Boolean products) and the fact that global morphisms respecting local divisibility entails local morphisms respecting the usual divisibility (a similar result for q. e. will be soon stated).

Idea of the proof

The model completeness of the theory T^* is established in the language of $\mathcal{L}_{\text{lor}} \cup \{\preceq, |_{\text{loc}}\}$ using Comer's result (a Feferman-Vaught result for Boolean products) and the fact that global morphisms respecting local divisibility entails local morphisms respecting the usual divisibility (a similar result for q. e. will be soon stated).

In order to prove the elimination of quantifiers of T^* , I will first describe the universal theory $T_{\forall}^*$ and secondly, prove that the amalgamation property for this theory $T_{\forall}^*$.

Theorem (G., 2025): The universal theory of T^* in the language $\mathcal{L}_{\text{lor}} \cup \{|\, , \preceq , |_{\text{loc}}\}$ is the theory of reduced f -rings satisfying the following axioms:

Theorem (G., 2025): The universal theory of T^* in the language $\mathcal{L}_{\text{lor}} \cup \{|\, , \preceq , |_{\text{loc}}\}$ is the theory of reduced f -rings satisfying the following axioms:

- the first convexity property: $\forall a \forall b (0 < a < b \rightarrow b \mid a),$

Theorem (G., 2025): The universal theory of T^* in the language $\mathcal{L}_{\text{lor}} \cup \{|\, , \preceq , |_{\text{loc}}\}$ is the theory of reduced f -rings satisfying the following axioms:

- the first convexity property: $\forall a \forall b (0 < a < b \rightarrow b \mid a)$,
- the glueing axioms for divisibility:
 $\forall a \forall b \forall c_1 \cdots \forall c_n ((bc_1 - a) \cdots (bc_n - a) = 0 \rightarrow b \mid a)$, for each $n \in \mathbb{N}$.

Theorem (G., 2025): The universal theory of T^* in the language $\mathcal{L}_{\text{lor}} \cup \{ |, \preceq, |_{\text{loc}} \}$ is the theory of reduced f -rings satisfying the following axioms:

- the first convexity property: $\forall a \forall b (0 < a < b \rightarrow b \mid a)$,
- the glueing axioms for divisibility:
 $\forall a \forall b \forall c_1 \cdots \forall c_n ((bc_1 - a) \cdots (bc_n - a) = 0 \rightarrow b \mid a)$, for each $n \in \mathbb{N}$.
- the local divisibility property: $\forall a \forall b \forall c (a \not\preceq bc - a \rightarrow b \mid_{\text{loc}} a)$.


Amalgamation property

Let $A, B, C \models T_{\forall}^*$ such that there exists $f: A \rightarrow B$ and $g: A \rightarrow C$ monomorphisms in the language $\mathcal{L}_{\text{lor}} \cup \{|\, \preceq, |_{\text{loc}}^{\text{m}}\}$.

Amalgamation property

Let $A, B, C \models T_{\forall}^*$ such that there exists $f: A \rightarrow B$ and $g: A \rightarrow C$ monomorphisms in the language $\mathcal{L}_{\text{lor}} \cup \{|\, \preceq, |\,^{\text{m}}_{\text{loc}}\}$.

We want to find $D \models T_{\forall}^*$ and monomorphisms $h: B \rightarrow D$ and $k: C \rightarrow D$ such that the following diagram is commutative:




Amalgamation property

By **Prestel-Schwartz**, there exists $\tilde{f}: \overline{\pi B}^{\text{con}} \rightarrow \overline{\pi A}^{\text{con}}$ and $\tilde{g}: \overline{\pi C}^{\text{con}} \rightarrow \overline{\pi A}^{\text{con}}$ continuous and surjective functions; that is:

$$\begin{array}{ccc} \overline{\pi B}^{\text{con}} & \xrightarrow{\tilde{f}} & \overline{\pi A}^{\text{con}} \\ & \nearrow \tilde{g} & \\ \overline{\pi C}^{\text{con}} & & \end{array}$$

Amalgamation property

By **Prestel-Schwartz**, there exists $\tilde{f}: \overline{\pi B}^{\text{con}} \rightarrow \overline{\pi A}^{\text{con}}$ and $\tilde{g}: \overline{\pi C}^{\text{con}} \rightarrow \overline{\pi A}^{\text{con}}$ continuous and surjective functions; that is:




This diagram can be completed using the coproduct of $\overline{\pi B}^{\text{con}}$ and $\overline{\pi C}^{\text{con}}$ over $\overline{\pi A}^{\text{con}}$, given by:

$$X = \overline{\pi B}^{\text{con}} \times_{\overline{\pi A}^{\text{con}}} \overline{\pi C}^{\text{con}} = \{(q_1, q_2) \in \overline{\pi B}^{\text{con}} \times \overline{\pi C}^{\text{con}} : \tilde{f}(q_1) = \tilde{g}(q_2)\}.$$


Amalgamation property and local morphisms

Thus we have the following commutative diagram:



Amalgamation property and local morphisms

Thus we have the following commutative diagram:



Theorem (G., 2025): Let $A \models T_{\forall}^*$ in the language $\mathcal{L}_{\text{lor}} \cup \{|\, \preceq, |\,^m_{\text{loc}}\}$ and let $B \models T^*$. Let $f: A \rightarrow B$ a monomorphism in the language $\mathcal{L}_{\text{lor}} \cup \{|\, \preceq, |\,^m_{\text{loc}}\}$. For $q \in \overline{\pi B}^{\text{con}}$ and $p \in \overline{\pi A}^{\text{con}}$ such $\tilde{f}(q) = f^{-1}(q) = p$, one has that $f_{pq}: A/p \rightarrow B/q$ is a monomorphism in the language $\mathcal{L}_{\text{or}} \cup \{|\, \}$.

Maximal local divisibility

In order to prove the previous theorem, the notion of local divisibility had to be strengthened to what I called a maximal local divisibility:

$$b \mid_{\text{loc}}^m a \leftrightarrow a = 0 \vee \exists e \left[e^2 = e \wedge ae \neq 0 \wedge e \preceq a \wedge b \mid ae \right. \\ \left. \wedge (a(1-e) = 0 \rightarrow b \mid a) \wedge (a(1-e) \neq 0 \rightarrow \right. \\ \left. \forall f (f^2 = f \wedge a(1-e)f \neq 0 \wedge f \preceq a(1-e) \rightarrow b \nmid a(1-e)f) \right].$$


Amalgamation property and local morphisms

Using the previous theorem for $\mathfrak{q} = (q_1, q_2) \in X$ and putting $p = \tilde{f}(q_1) = \tilde{g}(q_2) \in \overline{\pi A}^{\text{con}}$, one has monomorphisms $f_{pq_1}: A/p \rightarrow B/q_1$ and $g_{pq_2}: A/p \rightarrow C/q_2$ in the language $\mathcal{L}_{\text{or}} \cup \{|\}$

Amalgamation property and local morphisms

Using the previous theorem for $\mathfrak{q} = (q_1, q_2) \in X$ and putting $p = \tilde{f}(q_1) = \tilde{g}(q_2) \in \overline{\pi A}^{\text{con}}$, one has monomorphisms $f_{pq_1}: A/p \rightarrow B/q_1$ and $g_{pq_2}: A/p \rightarrow C/q_2$ in the language $\mathcal{L}_{\text{or}} \cup \{|\}$

By quantifier elimination of the theory RCVR, there exists a real closed valuation ring $R_{\mathfrak{q}}$ such that the following diagram is commutative:



Amalgamation property

Now consider the ring $D = \prod_{\mathfrak{q} \in X} R_{\mathfrak{q}}$, and let $h: B \rightarrow D$ and $k: C \rightarrow D$ be given by:

$$h(b) = (h_{\mathfrak{q}}(b + q_1))_{\mathfrak{q} \in X} \quad \text{and} \quad k(c) = (k_{\mathfrak{q}}(c + q_2))_{\mathfrak{q} \in X},$$

for $b \in B$ and $c \in C$, where $\mathfrak{q} = (q_1, q_2) \in X$.

Amalgamation property

Now consider the ring $D = \prod_{q \in X} R_q$, and let $h: B \rightarrow D$ and $k: C \rightarrow D$ be given by:

$$h(b) = (h_q(b + q_1))_{q \in X} \quad \text{and} \quad k(c) = (k_q(c + q_2))_{q \in X},$$

for $b \in B$ and $c \in C$, where $q = (q_1, q_2) \in X$.

It is rather technical to prove that h and k are monomorphisms in the language $\mathcal{L}_{\text{lor}} \cup \{|\, \preceq, |\,^{\text{m}}_{\text{loc}}\}$ and that $D \models T^*$.

Amalgamation property


Now consider the ring $D = \prod_{q \in X} R_q$, and let $h: B \rightarrow D$ and $k: C \rightarrow D$ be given by:

$$h(b) = (h_q(b + q_1))_{q \in X} \quad \text{and} \quad k(c) = (k_q(c + q_2))_{q \in X},$$

for $b \in B$ and $c \in C$, where $q = (q_1, q_2) \in X$.

It is rather technical to prove that h and k are monomorphisms in the language $\mathcal{L}_{\text{lor}} \cup \{|\cdot|, \preceq, |\cdot|_{\text{loc}}^{\text{m}}\}$ and that $D \models T^*$.

And finally the following diagram is commutative:



Two other cases

- Let $A = V_0 \times V_1$, where V_0, V_1 are two real closed valuation rings, then $T_2 = \text{Th}(A)$ is the theory of sc-regular projectable real closed rings with the first convexity property where the Boolean algebra of idempotents satisfies $\exists_{=4}$, i.e.: the formula saying that the structure has four elements.

Two other cases

- Let $A = V_0 \times V_1$, where V_0, V_1 are two real closed valuation rings, then $T_2 = \text{Th}(A)$ is the theory of sc-regular projectable real closed rings with the first convexity property where the Boolean algebra of idempotents satisfies $\exists_{=4}$, i.e.: the formula saying that the structure has four elements.

Using similar methods: it is easy to prove that T_2 admits elimination of quantifiers in the language $\mathcal{L}_{\text{lor}} \cup \{|\, \preceq, |\,_{\text{loc}}\}$

Two other cases

- Let $A = V_0 \times V_1$, where V_0, V_1 are two real closed valuation rings, then $T_2 = \text{Th}(A)$ is the theory of sc-regular projectable real closed rings with the first convexity property where the Boolean algebra of idempotents satisfies $\exists_{=4}$, i.e.: the formula saying that the structure has four elements.

Using similar methods: it is easy to prove that T_2 admits elimination of quantifiers in the language $\mathcal{L}_{\text{lor}} \cup \{ |, \preceq, |_{\text{loc}} \}$

- Now consider $F_2 = \text{Th}(F)$ where $F = F_1 \times F_2$ with F_1 and F_2 two real closed fields. In fact, F_2 is the theory of von Neumann regular real closed rings where there are only four idempotents.

Two other cases

- Let $A = V_0 \times V_1$, where V_0, V_1 are two real closed valuation rings, then $T_2 = \text{Th}(A)$ is the theory of sc-regular projectable real closed rings with the first convexity property where the Boolean algebra of idempotents satisfies $\exists_{=4}$, i.e.: the formula saying that the structure has four elements.

Using similar methods: it is easy to prove that T_2 admits elimination of quantifiers in the language $\mathcal{L}_{\text{lor}} \cup \{ |, \preceq, |_{\text{loc}} \}$

- Now consider $F_2 = \text{Th}(F)$ where $F = F_1 \times F_2$ with F_1 and F_2 two real closed fields. In fact, F_2 is the theory of von Neumann regular real closed rings where there are only four idempotents.

As the previous case, the theory F_2 has elimination of quantifiers in the language $\mathcal{L}_{\text{lor}} \cup \{ \preceq \}$, and so in the language $\mathcal{L}_{\text{lor}} \cup \{ |, \preceq, |_{\text{loc}} \}$.

The main theorem

Let A be a reduced projectable and divisible-projectable f -ring satisfying the first convexity property. Then A admits quantifier elimination in $\mathcal{L}_1 = \mathcal{L}_{\text{lor}} \cup \{ |, \preceq, |_{\text{loc}} \}$ if and only if one of the following conditions holds:

- (i) A is a real closed field,
- (ii) A is a product of two real closed fields,
- (iii) A is a von Neumann regular real closed ring without non-zero minimal idempotents,
- (iv) A is a real closed valuation ring,
- (v) A is a product of two real closed valuation rings,
- (vi) A is a sc-regular real closed ring without non-zero minimal idempotents.

The idea

The idea of the proof is to interpret the language $\mathcal{L}_1 = \mathcal{L}_{\text{lor}} \cup \{|\, \preceq, |\text{loc}\}$ in a previous language where I had proved the same result.

The idea

The idea of the proof is to interpret the language $\mathcal{L}_1 = \mathcal{L}_{\text{lor}} \cup \{|\, \preceq, |_{\text{loc}}\}$ in a previous language where I had proved the same result.

The previous language is $\mathcal{L}_2 = \mathcal{L}_{\text{lor}} \cup \{\text{div}(\cdot, \cdot)\}$, where

$$\text{div}(x, y) = c \iff c \preceq y \wedge \exists z \exists w (x = z + w \wedge z \perp w \wedge cy = z \wedge \forall w' (w' \neq 0 \wedge w' \perp (w - w') \rightarrow y \nmid w')).$$

Real closed domains

A domain D is **real closed** if the following conditions are verified:

- (i) $R = \text{qf}(D)$ is a real closed field,
- (ii) D is integrally closed (in R),
- (iii) $\forall a \forall b (0 < a < b \rightarrow b \mid a^2)$ is true in D .

Real closed domains

A domain D is **real closed** if the following conditions are verified:

- (i) $R = \text{qf}(D)$ is a real closed field,
- (ii) D is integrally closed (in R),
- (iii) $\forall a \forall b (0 < a < b \rightarrow b \mid a^2)$ is true in D .

Real closed domains are local rings, with m_D its unique maximal ideal.

Real closed domains

A domain D is **real closed** if the following conditions are verified:

- (i) $R = \text{qf}(D)$ is a real closed field,
- (ii) D is integrally closed (in R),
- (iii) $\forall a \forall b (0 < a < b \rightarrow b \mid a^2)$ is true in D .

Real closed domains are local rings, with m_D its unique maximal ideal.

The convex hull $\text{cH}(D, R)$ of D in R is a convex subring of R , therefore a real closed valuation ring of R with m_V its maximal ideal and $m_V \cap D = m_D$.

Real closed domains

A domain D is **real closed** if the following conditions are verified:

- (i) $R = \text{qf}(D)$ is a real closed field,
- (ii) D is integrally closed (in R),
- (iii) $\forall a \forall b (0 < a < b \rightarrow b \mid a^2)$ is true in D .

Real closed domains are local rings, with m_D its unique maximal ideal.

The convex hull $\text{cH}(D, R)$ of D in R is a convex subring of R , therefore a real closed valuation ring of R with m_V its maximal ideal and $m_V \cap D = m_D$.

Therefore $k_D = D/m_D$ is a subfield of $k_V = V/m_V$, and $\mathfrak{P}_D = (k_D, k_V)$ is a pair of real closed fields interpretable in the original structure D .

Real closed domains

A domain D is **real closed** if the following conditions are verified:

- (i) $R = \text{qf}(D)$ is a real closed field,
- (ii) D is integrally closed (in R),
- (iii) $\forall a \forall b (0 < a < b \rightarrow b \mid a^2)$ is true in D .

Real closed domains are local rings, with m_D its unique maximal ideal.

The convex hull $\text{cH}(D, R)$ of D in R is a convex subring of R , therefore a real closed valuation ring of R with m_V its maximal ideal and $m_V \cap D = m_D$.

Therefore $k_D = D/m_D$ is a subfield of $k_V = V/m_V$, and $\mathfrak{P}_D = (k_D, k_V)$ is a pair of real closed fields interpretable in the original structure D .

Proposition: If two real closed domains D_1 and D_2 are elementary equivalent, then $\mathfrak{P}_{D_1} \equiv \mathfrak{P}_{D_2}$.

Real closed domains

Is the converse true ?

Real closed domains

Is the converse true ?

What is $\text{tp}(x/A)$ in the $\text{Th}(D)$? (in order to carry on a back and forth between ω_1 -saturated structures).

Bibliography

1. G. Cherlin, M.A. Dickmann, *Real closed rings II. Model Theory*, Annals of Pure and Applied Logic 25 (1983), 213-231.
2. J. I. Guier, *Boolean products of real closed valuations rings and fields*, Annals of Pure and Applied Logic 112 (2001), 119-150.
3. J. I. Guier, *Projectable reduced f -rings admitting elimination of quantifiers*, Preprint.
4. J. I. Guier, *Elimination of quantifiers in a theory of real closed rings*, Annals of Pure and Applied Logic 176 (2025).
5. A. Prestel, J. Schmid, *Existentially closed domains with radical relations*, J. reine angew. Math. 407 (1990), 178-201.
6. A. Prestel, N. Schwartz, *Model theory of real closed rings*, In *Valuation theory and its applications*, vol. I (Saskatoon, SK, 1999), volume 32 of Fields Institute Communications, pp. 261-290, American Mathematical Society, Providence, RI, 2002.
7. N. Schwartz, *Real Closed Valuation Rings*, Communications in Algebra 37 (2009), pp 3796-3814.

Thanks

Thank you !